

**Multiparametric prostate MRI and cognitively targeted transperineal biopsy in patients with previous abdominoperineal resection and suspicion of prostate cancer.**

Authors:

Nienke L Hansen<sup>1,2</sup>, Iztok Caglic<sup>2,3</sup>, Laurence H Berman<sup>2,4</sup>, Christof Kastner<sup>2,5</sup>, Andrew Doble<sup>2,5</sup>, Tristan Barrett<sup>2,4</sup>

Affiliations:

1) Department of Diagnostic and Interventional Radiology, University Hospital RWTH Aachen, Aachen, Germany.

2) CamPARI Clinic, Addenbrooke's Hospital and University of Cambridge, Cambridge, UK

3) Department of Radiology, University Medical Centre Maribor, Maribor, Slovenia

4) Department of Radiology, Addenbrooke's Hospital and University of Cambridge, Cambridge, UK

5) Department of Urology, Addenbrooke's Hospital, Cambridge, UK

Corresponding address:

Tristan Barrett

Department of Radiology

University of Cambridge School of Clinical Medicine

Box 218, Cambridge Biomedical Campus

CB2 0QQ Cambridge, UK

Tristan.barrett@addenbrookes.nhs.uk

Tel: 01223 336895

Fax: 01223 330915

keywords: abdominoperineal resection, magnetic resonance imaging, prostate cancer, transperineal prostate biopsy

## **Acknowledgements**

Author 1 has received a research grant from RWTH Aachen University Hospital (Aachen, Germany). Author 4 is a lecturer for a 3-day satellite symposium organised by Siemens Healthcare and MedCom GmbH. Author 6 acknowledges support from Cancer Research UK, National Institute of Health Research Cambridge Biomedical Research Centre, Cancer Research UK and the Engineering and Physical Sciences Research Council Imaging Centre in Cambridge and Manchester and the Cambridge Experimental Cancer Medicine Centre. The Department of Urology has received sponsorship of various industry for organising Prostate MRI workshops.

## Introduction

Prostate cancer (PCa) is the commonest male cancer as well as the second leading cause of cancer death in men.<sup>1</sup> The standard cancer screening protocol includes serum prostate specific antigen (PSA) testing and digital rectal examination (DRE). If either are abnormal, prostate biopsy with subsequent histopathologic evaluation represents the gold standard for establishing the diagnosis.<sup>2</sup> Prostate biopsy can be performed either by the more traditional transrectal approach or by the increasingly popular transperineal route, using a transrectal ultrasound probe for needle guidance. In both methods, rectal access is essential. Abdominoperineal resection (APR) is a common surgical technique performed in a selected group of patients with inflammatory bowel disease (IBD), carcinoma of the distal rectum or anal canal and familial adenomatous polyposis (FAP). It involves resection of the distal part of the sigmoid colon, rectum, anus and the creation of a permanent end-colostomy.<sup>3</sup> As life expectancy in patients with IBD and FAP is not significantly lower than that of the general population and the 5-year-survival rate in stage T1 and T2 colorectal tumours treated with APR was found to be 78%, it is not uncommon to encounter post-APR patients presenting with a raised PSA and suspicion of prostate cancer.<sup>4</sup> The lack of rectal access means that such patients pose a diagnostic challenge.

Previously described diagnostic approaches in this cohort include biopsy by either a transperineal or transgluteal approach and under the guidance of ultrasound (transperineal, transabdominal or transurethral), magnetic resonance imaging (MRI) or computed tomography (CT).<sup>5-14</sup> Cancer detection rates ranged

from 40 - 82 %, <sup>7-14</sup> but even in the largest cohort, detection of intermediate to high grade prostate cancer (Gleason score  $\geq 7$ ) was low at 43%.<sup>12</sup> However, all studies to date have used a systematic biopsy approach and did not use MRI prior to biopsy to target suspicious areas.

Recently, multi-parametric Magnetic Resonance Imaging (mpMRI) has been shown to be effective for the detection and local staging of prostate cancer and can preferentially detect clinically relevant index tumours of higher grade and a size  $>5$  mm.<sup>15</sup> A normal mpMRI also has a high negative predictive ( $>90\%$ ) for the presence of clinically significant disease.<sup>16</sup> Using transperineal MR/US fusion biopsy, the detection rate for clinically significant disease has been shown to be as high as 72%.<sup>17</sup> The aim of this study was to evaluate if the use of multiparametric MRI and subsequent cognitively targeted US-guided (by visual registration) transperineal biopsy improves cancer detection and particularly detection of significant cancer in patients with abdominoperineal resection and a suspicion of prostate cancer.

## **Patients and Methods**

### **Study population**

This single-institution retrospective study was part of a service evaluation of transperineal prostate biopsies with the need for informed consent for data analysis waived by the local ethics committee (Local audit evaluation number 4058). The study population consisted of 11 consecutive patients who presented to our institution between March 2010 and May 2015 with elevated PSA, a history of abdominoperineal resection of the rectum and undergoing prostate MRI at our institution prior to biopsy.

### **Magnetic resonance imaging**

Patients underwent prostate MRI on a 1.5T MR450 or 3.0T Discovery MR750 HDx (GE Healthcare, Waukesha, USA) with an 8-16 channel surface phased array coil. Axial Fast Spin Echo T1-weighted images of the pelvis, along with T2-Weighted Fast Recovery Fast Spin Echo images of the prostate were acquired in the axial, sagittal and coronal planes (TE/TR = 85/3700-5000 ms; field-of-view (FOV) 24x24 cm; matrix 256x256; slice thickness 3 mm; gap 1 mm). Diffusion-weighted (DW) imaging was performed using a spin-echo echo-planar imaging pulse sequence (TE/TR=60/3000-3400 ms; matrix 256x256; slice thickness 4 mm; gap 0 mm, FOV: 1.5T: 24x24 cm, 3.0T: 28x28 cm; parallel imaging factor of 2; signal averages: 3 for 1.5T and 8 for 3.0T. The following b-values were

acquired: b-150, b-750, b-1,400 s/mm<sup>2</sup>; apparent diffusion coefficient (ADC) maps were automatically calculated and stored.

### Image analysis

The MR images were prospectively reported by body radiologists experienced in reading prostate MRI. T2WI and DWI sequences were evaluated using a Likert scale of tumour probability, based on the Prostate Imaging Reporting and Data System (PI-RADS) structured scoring criteria developed by the European Society of Urogenital Radiology (ESUR).<sup>18</sup> The Likert-based scoring system was as follows: 1 = cancer highly unlikely, 2 = cancer unlikely, 3 = indeterminate for cancer, 4 = cancer likely, 5 = cancer highly likely. A positive MRI was defined as a score of  $\geq 3$ . The location of the lesion was communicated in the report descriptively for earlier cases, and subsequently using the standardized sector map first proposed by Dickinson et al.<sup>19</sup>

### Biopsy

Patients with a Likert score of 3-5 underwent transperineal ultrasound-guided biopsies under general anaesthesia. Transperineal biopsies were performed by a urologist (XX) with 29 years experience in prostate biopsy and 8 years experience in transperineal biopsy. The procedure was performed using transperineal ultrasound-guidance from one of two experienced radiologists (XX, XX). Each patient was placed in the lithotomy position, with the scrotum retracted

anteriorly. Bladder filling was not routinely employed as a transvesicular imaging plane was not chosen.

All ultrasound scanning was performed on a Toshiba Aplio 500 machine (Toshiba Medical Systems, UK) using a curved linear array ultrasound transducer with a centre frequency of 3.5MHz, range 1.9-6 MHz. The transducer was enclosed in a rubber sheath with sterilised ultrasound gel for transmission. Direct contact was made to the skin in the midline sagittal plane, to allow lateral transperineal needle access (Figure 1). Biopsies were obtained using an automatic spring driven biopsy device mounted with an 18-gauge biopsy needle (Bard biopsy systems, Tempe, USA) via a caudal approach to the prostate. The needle was directed to target areas by visual registration and systematically to background prostate by the operator, with position confirmed on US prior to sampling (Figure 2).

### Histopathology

Biopsies were reviewed by a single sub-specialist uropathologist. Each core was measured, and the total number of cores with cancer and the percentage of cancer recorded. Pathology reports identified the Gleason Score for each side of the prostate, and the target.

## Results

11 men with a median age of 68 years (range 63 - 77 years) were included with a median PSA of 9.4 ng/ml (mean 13.6; range 3.2 - 60 ng/mL); **Table 1**. The mean prostate volume was 37.8 ml (median 35; range 24 - 49 ml). Abdominoperineal resection had been performed for ulcerative colitis (7 patients), Crohn's disease, Hereditary nonpolyposis colorectal cancer, rectal carcinoma, and Canada-Cronkheit syndrome. The mean interval from MRI to biopsy was 57.6 days (range 22 - 106 days).

Transperineal ultrasound guided biopsies were successfully completed in 9 cases. On average 16.7 (range 11 - 27) biopsy cores with a median of 5 (range 3-7) target cores were obtained under ultrasound guidance. All patients undergoing biopsy were discharged on the day of procedure, 2/9 patients experienced acute urinary retention after biopsy, 1 described mild perineal discomfort, the other 6 had no ill effects. Of the 9 biopsies performed, 7 (78%) revealed prostate cancer with a Gleason score ranging from 6 - 10. 6/9 (67%) cases revealed a Gleason score of  $\geq 7$ , 1/9 cases (11%) revealed a Gleason score of 6. Median cancer core length was 2 mm (range 1 - 19mm). Treatment choices in patients with confirmed prostate cancer included open radical prostatectomy in 2 patients, external beam radiation therapy in 2, active surveillance in 2, and androgen deprivation therapy in 1.



9 patients exhibited suspicious lesions on MRI (**Figure 2**). Five patients had suspicious lesions in the posterior peripheral zone and four patients in the anterior transition zone. The median lesion diameter was 15 mm (range 7-25 mm). All 7 patients with highly suspicious lesions (Likert 4-5) on MRI had subsequent biopsy-proven tumour: 6 cases revealed Gleason score  $\geq 7$  cancer, with 1 case of Gleason grade 3+3. Biopsies from the target and adjacent area did more often contain cancer than background cores with higher number of positive cores in all seven cases, higher percentages of cancer involvement in all seven cases and/or higher Gleason scores in five cases.

The 2 cases with MR lesions of an intermediate probability score of 3/5 revealed benign tissue at biopsy. Two patients with a negative MRI did not undergo biopsy. The 4 patients with no histological diagnosis of prostate cancer had stable PSA levels at follow-up, with PSA-monitoring performed at 3 months for one year and 6-monthly thereafter; mean follow-up 17.5 months (range 7 - 28 months)

## Comment

Our study shows that multiparametric prostate MRI is feasible and useful for targeted transperineal biopsy guided by visual registration in patients with abdominoperineal resection and a suspicion of prostate cancer. The overall cancer detection rate was 7/9 (78%) with 6/9 (67%) cases demonstrating a Gleason score of 7 or higher. This compares favourably to previously published studies without prior MRI, reporting overall cancer detection rates of 20% (2 of 10 patients), 40% (2/5), 43% (3/7), 60% (6/10) and 82 % (23/28).<sup>7-10,12</sup> It has previously been shown that pre-biopsy MR imaging increases biopsy performance in detecting prostate cancer and especially clinically significant cancer, via both a transperineal and transrectal approach using MR/US fusion techniques.<sup>17,20</sup> Although our high detection rate of clinically significant cancers may partially be explained by patients without rectal access presenting with larger and more significant cancers due a delayed diagnosis,<sup>20</sup> our detection rate of 67% for Gleason score  $\geq 7$  disease is notably higher than any of the equivalent series that have previously been reported.

Our data suggests that patients with abdominoperineal resection benefit from a pre-biopsy multiparametric MRI and, in the case of suspicious lesions with a Likert score of 4-5, should undergo targeted transperineal biopsy guided by visual registration. In our study population, 6/7 cases with highly suspicious MRI lesions (Likert 4-5) revealed clinically significant adenocarcinomas with a

Gleason score of 7 or higher, with 1 case of low-risk Gleason score 6 prostate cancer. The results are in accordance with a meta-analysis of patients with rectal access undergoing MRI prior to initial targeted biopsy of the prostate with an overall detection rate of 65% for prostate cancer and 49% for significant disease.<sup>21</sup> Additionally, a recent study using targeted transperineal biopsy with rectal access for MR/US fusion found that 85% (89/104) of patients with highly suspicious lesions on MRI were diagnosed with prostate cancer in the biopsy, including 60% (62/104) with high risk cancer of GS of 7 or greater.<sup>17</sup> A fusion approach is not possible in the setting of prior APR and image guidance by visual registration despite its operator-dependence is the only means of biopsy targeting. Previous studies have reported contradicting results when comparing visually-registered and software-registered MR/US fusion targeted transrectal biopsies after prebiopsy MRI: with one demonstrating equivalence,<sup>20</sup> another showing increased detection with software-registration biopsy, especially for smaller lesions,<sup>22</sup> and a third, larger study, concluding that software-registration biopsy was more histologically informative but did not increase cancer detection.<sup>23</sup> An alternative to biopsies guided by visual estimation would be real-time in-bore MRI-targeted biopsies, via a transperineal or transgluteal approach,<sup>24,25</sup> however, this is cost- and time-intensive, requires special MR-safe equipment, and is not available at all centers. In our study, the 2 cases with Likert 3 MR lesions revealed benign tissue upon biopsy. It may therefore be possible to avoid biopsy in patients with intermediate risk MRI, but this is a controversial area. Previous studies in patients without APR, showed a yield of

any cancer of 20-25% in intermediate probability cases, although typically the majority of these are low risk cancers.<sup>26,27</sup>

A limitation of our study is its retrospective nature and the small population size, especially the patients without suspicious lesions on MRI. The lack of prostatectomy as a gold standard in all but two patients is also a limitation, and we therefore cannot exclude some smaller tumours being missed. Another limitation of MRI –targeted transperineal biopsies guided by visual registration is the difficulty of ultrasound guidance. Needle visualisation can be challenging due to lateral entry point of the needle to the ultrasound probe and the off-plane trajectory taken, additionally the beam may be scattered by the fibromuscular layers of the pelvic floor or surgery-related scarring. Theoretically these effects could be exacerbated in glands with larger volume, more heterogeneous transition zone, however, significant BPH was not present in our cohort (gland volume 24 - 49 ml). Patient body mass index (BMI) is less likely to have an affect given the transperineal approach. Steps taken to improve visualisation included small movements of the needle to ascertain position indirectly through tissue motion, incremental tilting and rotation of the probe to change beam position, or even switching to the transverse plane (**Figure 3**). Although not employed here, consideration could also be given to catheterisation to aid urethral land-marking and help establish the prostatic midpoint.

We describe the first use of MRI to aid targeting of prostate biopsies in patients

post APR, with the approach achieving promising results in this challenging patient group. Our findings could also be extended to the group of patients with ileal pouch/anal anastomosis, where transpouch biopsy of the prostate may risk pouch fistulas or abscesses.<sup>28-30</sup>

## **Conclusions**

In conclusion, the use of multiparametric prostate MRI and subsequent targeted transperineal biopsy guided by visual registration can aid in the diagnostic pathway of patients with abdominoperineal resection and a suspicion of prostate cancer.

## References

1. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin* 2015.
2. Heidenreich A, Bastian PJ, Bellmunt J, et al. European Association of Urology. EAU guidelines on prostate cancer. part 1: screening, diagnosis, and local treatment with curative intent-update 2013. *Eur Urol*. 2014 Jan;65(1):124-37.
3. Perry WB, Connaughton JC. Abdominoperineal Resection: How Is It Done and What Are the Results? *Clin Colon Rectal Surg*. 2007 Aug; 20(3): 213–220.
4. Olsheski M, Schwartz D, Rineer J, et al. A population-based comparison of overall and disease-specific survival following local excision or abdominoperineal resection for stage I rectal adenocarcinoma. *J Gastrointest Cancer*. 2013 Sep;44(3):305-12.
5. Rathaus V, Richter S, Nissenkorn I, et al. Transperineal ultrasound examination in the evaluation of prostatic size. *Clin Radiol* 1991 ;44:383-385.
6. Krauss DJ, Clark KG, Nsouli IS, et al. Prostate biopsy in patients after proctectomy. *J Urol*. 1993 Mar;149(3):604-6.
7. Twidwell JJ, Matthews RD, Huisam TK, et al. Ultrasound evaluation of the prostate after abdominoperineal resection. *J Urol*. 1993 Sep;150(3):902-4.

8. Seaman EK, Sawczuk IS, Fatal M, et al. Transperineal prostate needle biopsy guided by transurethral ultrasound in patients without a rectum. *Urology* 1996; 47(3):353-5.
9. Filderman PS, Jacobs SC. Prostatic ultrasound in the patient without a rectum. *Urology*. 1994; 43(5):722-4.
10. Papanicolaou N1, Eisenberg PJ, Silverman SG, et al. Prostatic biopsy after proctocolectomy: a transgluteal, CT-guided approach. *AJR Am J Roentgenol*. 1996 Jun;166(6):1332-4.
11. D'Amico AV, Tempany CM, Cormack R, et al. Transperineal magnetic resonance image guided prostate biopsy. *J Urol*. 2000 Aug;164(2):385-7.
12. Shinohara K, Gulati M, Koppie T, et al. Transperineal prostate biopsy after abdominoperineal resection. *J Urol*. 2003; 169(1):141-4.
13. Cantwell CP, Hahn PF, Gervais DA, et al. Prostate biopsy after ano-rectal resection: value of CT-guided trans-gluteal biopsy. *Eur Radiol*. 2008 Apr;18(4):738-42.
14. Goenka AH, Remer EM, Veniero JC, et al. CT-Guided Transgluteal Biopsy for Systematic Random Sampling of the Prostate in Patients Without Rectal Access. *AJR Am J Roentgenol*. 2015 Sep;205(3):578-83.
15. Turkbey B, Mani H, Shah V, et al. Multiparametric 3T Prostate Magnetic Resonance Imaging to Detect Cancer: Histopathological Correlation Using prostatectomy specimens processed in customized magnetic resonance imaging based molds. *J Urol*. 2011 Nov;186(5):1818-24.


16. Yerram NK, Volkin D, Turkbey B, et al. Low suspicion lesions on multiparametric magnetic resonance imaging predict for the absence of high-risk prostate cancer. *BJU Int.* 2012 Dec;110(11 Pt B):E783-8.
17. Kuru TH, Roethke MC, Seidenader J, et al. Critical evaluation of magnetic resonance imaging targeted, transrectal ultrasound guided transperineal fusion biopsy for detection of prostate cancer. *J Urol.* 2013 Oct;190(4):1380-6.
18. Barentsz JO, Richenberg J, Clements R, et al. ESUR prostate MR guidelines 2012. *Eur Radiol Eur Radiol* 2012; 22: 746–757.
19. Dickinson L, Ahmed HU, Allen C, et al. Magnetic resonance imaging for the detection, localisation, and characterisation of prostate cancer: recommendations from a European Consensus Meeting. *Eur Urol* 2011;59:477e94.
20. Umbreit EC, Dozois EJ, Crispen PL, et al. Radical prostatectomy for prostate cancer after ileal pouch-anal anastomosis offers oncologic control and sustains quality of life. *J Am Coll Surg.* 2010 Feb;210(2):232-9.
21. Schoots IG, Roobol MJ, Nieboer D, et al. Magnetic resonance imaging-targeted biopsy may enhance the diagnostic accuracy of significant prostate cancer detection compared to standard transrectal ultrasound-guided biopsy: a systematic review and meta-analysis. *Eur Urol.* 2015; 68(3):438-50.
22. Puech P, Rouviere O, Renard-Penna R, et al. Prostate cancer diagnosis: multiparametric MR-targeted biopsy with cognitive and transrectal US-MR



- fusion guidance vs. Systematic biopsy- prospective multicenter study. Radiology 2013; 268:461-469.
23. Cool DW, Zhang X, Romagnoli C, et al. Evaluation of MRI-TRUS fusion versus cognitive registration accuracy for MRI-targeted, TRUS-guided prostate biopsy. AJR Am J Roentgenol. 2015 Jan;204(1):83-91.
24. Wysock JS, Rosenkrantz AB, Huang WC, et al. A prospective, blinded comparison of magnetic resonance (MR) imaging-ultrasound fusion and visual estimation in the performance of MR-targeted prostate biopsy: the PROFUS trial. Eur Urol. 2014 Aug;66(2):343-51.
25. Bodelle B, Naguib NN, Schulz B, et al. 1.5-T magnetic resonance-guided transgluteal biopsies of the prostate in patients with clinically suspected prostate cancer: technique and feasibility. Invest Radiol. 2013 Jun;48(6):458-63.
26. Gaziev G, Wadhwa K, Barrett T, et al. Defining the learning curve for multiparametric magnetic resonance imaging (MRI) of the prostate using MRI-transrectal ultrasonography (TRUS) fusion-guided transperineal prostate biopsies as a validation tool. BJU Int. 2014 Aug 7.
27. Renard-Penna R, Mozer P, Cornud F, et al. Prostate Imaging Reporting and Data System and Likert Scoring System: Multiparametric MR Imaging Validation Study to Screen Patients for Initial Biopsy. Radiology. 2015 May;275(2):458-68.
28. Kani HT, Shen B. Male issues of the ileal pouch. Inflamm Bowel Dis. 2015; 21(3):716-22.


29. Fergany AF, Angermeier KW. A technique of transrectal ultrasound guided transperineal random biopsy in patients with ulcerative colitis and ileal pouch. *J Urol* 2000; 163:205-6.
30. Shen B, Angermeier KW, Remzi FH, et al. Screening and diagnosis of prostate cancer in patients with ileal pouch-anal anastomosis: consensus from an expert panel. *Am J Gastroenterol*. 2011; 106(2):186-9.

## Legends to illustrations




**Fig. 1 Transperineal ultrasound views**

A: ultrasound image in the sagittal (longitudinal) plane. B: sagittal T2-weighted MR image at the same level. PZ = peripheral zone of prostate; TZ = transition zone; CS = corpus spongiosum; BI = bladder; P = pubic bone



**Fig. 2 High probability MRI target**

66 year old patient with PSA 11.74 ng/ml and previous total colectomy for Canada-Cronkhite syndrome. A-C: MRI shows a high probability 2.5 cm lesion in the anterior right base transition zone (\*) with low T2 signal intensity (A) and restricted diffusion with high signal on the b-1400 images (B) and low ADC value (C). D: Sagittal plane ultrasound image of transperineal biopsy needle (arrow) guided to this region. Subsequent pathology confirmed Gleason 4+3 disease



**Fig. 3 Ultrasound biopsy guidance: Switching to coronal plane for needle visualisation**

68 year old patient with PSA 3.2 ng/ml and strong family history of prostate cancer and low probability MRI. A: Ultrasound switched to coronal (transverse) plane to aid needle visualisation (arrow). B: Coronal T2-weighted MR image at an equivalent position

## Tables

**Table 1** Results for patients who underwent transperineal prostate biopsy listed in order of date of procedure

| Patient | Age (years) | Indication for anorectal resection | Time since resection (years) | PSA (ng/mL) | Volume (mL) | PSA Density (ng/mL/mL) | MRI results | Max Lesion diameter (mm) | Interval MRI to biopsy (days) | Number of Cores (n) | Number of positive Cores (n) | CCL (mm) | Biopsy Result (Gleason score) | Management                |
|---------|-------------|------------------------------------|------------------------------|-------------|-------------|------------------------|-------------|--------------------------|-------------------------------|---------------------|------------------------------|----------|-------------------------------|---------------------------|
| 1       | 72          | UC                                 | 22                           | 60          | 28          | 2.14                   | +           | 15                       | 22                            | 11                  | 3                            | 3        | 5+5                           | ADT                       |
| 2       | 77          | Rectal Ca                          | 5                            | 15.7        | 31          | 0.51                   | +           | 16                       | 70                            | 19                  | 6                            | 10       | 3+5                           | EBRT                      |
| 3       | 77          | UC                                 | 2                            | 14.8        | 45          | 0.33                   | +           | 14                       | 34                            | 13                  | 1                            | 1        | 4+3                           | Neoadjuvant ADT, EBRT     |
| 4       | 65          | UC                                 | 43                           | 5.2         | 24          | 0.22                   | +           | 10                       | 106                           | 12                  | 2                            | 1        | 3+3                           | AS                        |
| 5       | 65          | UC                                 | n/a                          | 7.1         | 26          | 0.27                   | +           | 9                        | 58                            | 27                  | 0                            |          | Benign                        | PSA Follow-up, stable     |
| 6       | 66          | CCS                                | 30                           | 11.7        | 34          | 0.35                   | +           | 25                       | 83                            | 16                  | 5                            | 2        | 3+4                           | AS                        |
| 7       | 63          | HNPCC                              | 14                           | 3.2         | 49          | 0.07                   | +           | 7                        | 68                            | 21                  | 0                            |          | Benign                        | PSA Follow-up, stable     |
| 8       | 70          | UC                                 | 7                            | 9.4         | 44          | 0.21                   | +           | 15                       | 43                            | 13                  | 5                            | 2        | 4+5                           | Neoadj ADT, prostatectomy |
| 9       | 65          | UC                                 | 25                           | 3.9         | 35          | 0.11                   | +           | 16                       | 34                            | 18                  | 12                           | 19       | 4+5                           | prostatectomy             |
| 10      | 68          | Crohn's                            | 40                           | 8.8         | 59          | 0.15                   | -           |                          |                               |                     |                              |          |                               | PSA Follow-up, stable     |
| 11      | 68          | UC                                 | 6                            | 9.6         | 41          | 0.23                   | -           |                          |                               |                     |                              |          |                               | PSA Follow-up, stable     |

HNPCC = Hereditary nonpolyposis colorectal cancer, UC = ulcerative colitis, CCS = Canada-Cronkheit syndrome, AS = active surveillance, ADT = androgen deprivation therapy, GI = Gleason, EBRT = external beam radiotherapy, PSA = prostate specific antigen

